

Nematic Colloidal Crystals and Superstructures

I. Musevic^{1,2}

¹*J. Stefan Institute, Jamova 39, SI-1000 Ljubljana, Slovenia*

²*Faculty of Mathematics and Physics, University of Ljubljana, Jadraska 19, SI-1000, Ljubljana, Slovenia*

1. INTRODUCTION

We shall discuss the general properties and recent development in the field of nematic colloids, which are dispersions of microparticles in a nematic liquid crystal. As the nematic liquid crystal is an orientationally ordered fluid, it interacts with the surfaces of inclusions that impose a preferential order at a curved interface. This interaction generates topological defects, which are responsible for structural forces between dispersed microparticles [1].

2. STRUCTURAL FORCES AND COLLOIDAL ASSEMBLY IN NEMATIC LIQUID CRYSTALS

The structural forces in liquid crystals are extremely strong, anisotropic and long-range, and are responsible for directed assembly of a broad variety of colloidal superstructures in liquid crystals: 2D colloidal crystals [2], colloidal wires, assembled by entangled topological defects [3], superstructures in the mixtures of large and small colloidal particles [4] and a broad variety of 2D nematic colloidal crystals [5]. In all cases, the colloidal binding energy is several orders of magnitude stronger compared to water based colloids, and could provide assembly of nanometer-sized colloidal particles of various shapes [6]. The methods of manipulation of microparticles and the mechanisms of their assembly in the nematic liquid crystal will be discussed. Comparison between theoretical predictions based on Landau-de Gennes theory and the experiments will be addressed. Possible application of directed and robust colloidal assembly for photonic devices will be discussed, as it has recently been shown, that a single nematic microdroplet in a liquid or polymer matrix provides a novel class of tunable optical microresonators. Light can be trapped in a microdroplet due to the total internal reflection and the resonant eigenmodes can be tuned by an external electric field [7]. This could be a basic tunable optical resonant element for future technological platform for soft matter photonic devices.

ACKNOWLEDGEMENT

The author would like to thank Miha Škarabot for his valuable contribution to this work. Contributions of Miha Ravnik, Slobodan Žumer, Uroš Tkalec, Igor Poberaj, Dušan Babič, Natan Osterman, Andriy Nych, Ulyana Ognysta and Vassili Nazarenko are also acknowledged.

REFERENCES

1. P. Poulin, H. Stark, T. C. Lubensky, and D. A. Weitz, *Science* **275**, 1770(1997).
2. I. Muševič, M. Škarabot, U. Tkalec, M. Ravnik, S. Žumer, *Science* **313**, 954(2006)
3. M. Ravnik, M. Škarabot, S. Žumer, U. Tkalec, I. Poberaj, D. Babič, N. Osterman, I. Muševič, *Phys. Rev. Lett.* **99**, 247801(2007).
4. M. Škarabot, M. Ravnik, S. Žumer, U. Tkalec, I. Poberaj, D. Babič, I. Muševič, *Phys. Rev. E* **77**, 061706. (2008).
5. U. Ognysta, A. Nych, V. Nazarenko, I. Muševič, M. Škarabot, M. Ravnik, S. Žumer, I. Poberaj, D. Babič, *Phys. Rev. Lett.* **100**, 217803(2008).
6. F. R. Hung, *Phys. Rev. E* **79**, 021705(2009).
7. M. Humar, M. Ravnik, S. Pajk, I. Muševič, *Nature Photon.* **3**, 595(2009).